

IN THE CLAIMS

Please amend the claims as follows:

1. (Currently Amended) A method of stimulating a HIV1-specific CD8⁺ response in a human infected with an HIV retrovirus said method comprising:
administering to the human, a recombinant virus, which enters the cells of the human and intracellularly produces HIV specific peptides for presentation on the cell's MHC class I molecules,
where said peptides are presented in an amount sufficient to stimulate a protective CD8⁺ HIV structural antigen response, and
where said human
 - i. has a viral load of less than 10,000 viral copies per ml of plasma and a CD4⁺ cell count of above 500 cells/ml, and
 - ii. has been treated with one or more anti-viral agents, which contributed to a lower viral copy and higher CD4⁺ cell count than before treatmentwhere said HIV specific peptides comprise HIV Gag, Gp120, Nef or Pol peptides.
2. (Previously Presented) A method of claim 1 wherein the human has been treated with anti-viral agents, which resulted in the human having a viral load of less than 1,000 viral copies per ml of blood serum and a CD4⁺ cell count of above 500 cells/ml.
3. (Original) A method of claim 2 wherein the anti-viral agents comprise a combination of protease inhibitors and inhibitors of reverse transcriptase.
4. (Canceled).
5. (Previously Presented) A method of claim 1 wherein the recombinant virus is an attenuated recombinant virus.

-
6. (Previously Presented) A method of claim 5 wherein the attenuated recombinant virus comprises a pox virus.
 7. (Previously Presented) A method of claim 6 wherein the attenuated recombinant pox virus comprises NYVAC or ALVAC.
 8. (Previously Presented) A method of claim 6 wherein the recombinant pox virus comprises MVA.
 9. (Original) A method of claim 1 where the vaccine is administered a second time.
 10. (Previously Presented) A method of claim 1 wherein the HIV specific peptides are structural viral peptides.
 11. (Canceled).
 12. (Original) A method of claim 1 wherein the vaccine further comprises an adjuvant.
 13. (Original) A method of claim 1 further comprising administering interleukin 2 or CD40 ligand in an amount sufficient to potentiate the CD8⁺ response.
 14. (Previously Presented) A method of claim 1 where the human has been infected with HIV and has demonstrated repeated and sustained proliferative T-cell responses to Gp120 envelope protein.
 15. (Previously Presented) A method of claim 14 where the human has demonstrated repeated and sustained proliferative T-cell responses to p24 Gag antigen.
 16. (Previously Presented) A method of claim 1 where the human is infected with HIV and is further tested by a skin test for a hypersensitive response to p24 Gag antigen.

17. (Previously Presented) A method of claim 1 where the human is infected with HIV and is further tested by a skin test for a hypersensitive response to Gp120 envelope antigen.

18. (Currently Amended) A method of maintaining a reduced ~~reducing~~ viral load in a mammal infected with an immunodeficiency retrovirus said method comprising:

administering to the mammal a recombinant virus, which enters the cells of the mammal and intracellularly produces immunodeficiency retroviral specific peptides for presentation on the cell's MHC class I molecules,

where said peptides are presented in an amount sufficient to stimulate a protective CD8⁺ HIV ~~structural~~ antigen response and thereby maintain a reduced viral load in the mammal, and where said mammal

i. has an immunodeficiency retroviral load of less than 10,000 viral copies per ml of plasma and a CD4⁺ cell count of above 500 cells/ml prior to administration of the recombinant virus, and

ii. has been treated with one or more anti-viral agents, which contributed to a lower viral copy and higher CD4⁺ cell count before treatment where said peptides comprise immunodeficiency retroviral Gag, Gp120, Nef or Pol peptides.

19. (New) A method of stimulating a HIV1-specific CD8⁺ response in a human infected with an HIV retrovirus said method comprising:

administering to the human, a recombinant virus, which enters the cells of the human and intracellularly produces HIV specific peptides for presentation on the cell's MHC class I molecules,

where said peptides are presented in an amount sufficient to stimulate a protective CD8⁺ HIV antigen response, and

where said human

i. has a viral load of less than 10,000 viral copies per ml of plasma and a CD4⁺ cell count of above 500 cells/ml, and

ii. has been treated with one or more anti-viral agents, which contributed to a lower viral copy and higher CD4⁺ cell count than before treatment.

20. (New) A method of stimulating a HIV1-specific CD8⁺ response in a human infected with an HIV retrovirus said method comprising:

administering to the human, a recombinant virus, which enters the cells of the human and intracellularly produces HIV specific peptides for presentation on the cell's MHC class I molecules,

where said peptides are presented in an amount sufficient to stimulate a protective CD8⁺ HIV structural antigen response, and

where said human

i. has a viral load of less than 10,000 viral copies per ml of plasma and a CD4⁺ cell count of above 500 cells/ml, and

ii. has been treated with one or more anti-viral agents, which contributed to a lower viral copy and higher CD4⁺ cell count than before treatment

where said HIV specific peptides comprise Gag, Pol, Env peptides or a combination thereof.